

Dutt and Stead from the United States recommend a regimen consisting of rifampin (600 mg) and isoniazid (300 mg) given daily for about two months, followed by rifampin (600 mg) and isoniazid 15 mg per kg of body weight) daily or twice a week (depending on patient reliability) for another seven months. They add ethambutol hydrochloride (15 mg per kg) during the first two months if a patient has shown high initial drug resistance.

Many centers now recommend without reservation short-course chemotherapy for the initial treatment of genitourinary tuberculosis. The results may be even better than in pulmonary tuberculosis because fewer bacilli are usually found in renal lesions and rifampin is highly concentrated in the urine.

STUART D. BOYD, MD

REFERENCES

- Dutt AK, Stead WW: Chemotherapy of tuberculosis for the 1980's. *Clin Chest Med* 1980 May; 1(2):243-252
 Farer LS, Long MW, Snider DE Jr, et al: Correspondence. *Am Rev Respir Dis* 1981 Nov; 124:657
 Gow JG: Genitourinary tuberculosis: A 7-year review. *Br J Urol* 1979 Aug; 51:239-244

Single-Dose Therapy for Lower Urinary Tract Infection

ACUTE CYSTITIS is one of the two most frequent indications for administering antimicrobial therapy in adult women. Recent studies have shown that in non-pregnant adult women single-dose antimicrobial therapy may be as effective as conventional 7- to 14-day regimens.

Controlled trials have shown that amoxicillin trihydrate, sulfisoxazole and trimethoprim-sulfamethoxazole are effective when given in single-dose regimens in this population. In contrast, cephaloridine, cefaclor and cephalexin monohydrate have been shown to be poor choices. Caution should be exercised in extrapolating data to other agents without appropriate clinical evaluation. Likewise, single-dose regimens have been less effective than conventional treatment in pregnant women and children and should not be used in these patients.

Potential benefits of single-dose therapy include reduced medication costs, improved patient compliance with drug regimens, fewer side effects and a reduction in the rate of emergence of resistant bacteria. Additionally, the failure of a single-dose regimen to eradicate bacteriuria (when the organism is sensitive) may indicate more invasive infections and, some suggest, a need for closer follow-up and more extensive investigation to identify structural or functional abnormalities of the urinary tract. Careful bacteriologic follow-up evaluation should be done irrespective of the duration of therapy.

Although information regarding optimal choice and dose of an antimicrobial agent for single-dose treatment remains to be refined, oral amoxicillin (2.0 grams), sulfisoxazole (1.0 grams) or trimethoprim-sulfamethoxazole (160 and 800 mg, respectively) appear to be rational choices.

Administration of single-dose antibiotic drugs for lower urinary tract infection is safe and effective therapy for nonpregnant adult women.

PAUL F. SOUNEY

REFERENCES

- Harris RE, Gilstrap LC III, Pretty A: Single-dose antimicrobial therapy for asymptomatic bacteriuria during pregnancy. *Obstet Gynecol* 1982 May; 59:546-549
 Kunin CM: Duration of treatment of urinary tract infections. *Am J Med* 1981 Nov; 71:849-854
 Souney P, Polk BF: Single-dose antimicrobial therapy for urinary tract infections in women. *Rev Inf Dis* 1982 Jan-Feb; 4:29-34

Hospital Admission Rates in Men Who Have Had Vasectomies

FEARS based on findings from studies in animals about the long-term health of men who have had vasectomies have not been borne out in a large follow-up study. Using the records of a large health maintenance organization, the incidence of first hospital admission after vasectomy in 6,092 men (20,491 man-years of follow-up) was evaluated. Hospital admissions for various classes of disease and for a variety of specific illnesses were compared with the rates in members of the same health maintenance organization who had not had vasectomies. There was no evidence of an increased prevalence of arteriosclerotic disease in the men who had had vasectomies, the principal concern raised by the experiments in animals. The highest standardized rate ratio (vasectomized to nonvasectomized) for first hospital admissions was for diseases of the genitourinary system, whose diagnosis may have been prompted by the vasectomy.

These findings were further evaluated according to time since the vasectomy because it was assumed that hospital admittance causally associated with vasectomy would either increase or decrease in frequency in some regular fashion with time. No specific disease entity followed this pattern, with the exception of hospital admissions for mental disorders, which were less frequent in men who had had vasectomies than in men who had not and continued to decline with time after the vasectomy procedure.

ALEXANDER M. WALKER, MD, DrPH

REFERENCES

- Clarkson TB, Alexander NJ: Long-term vasectomy: Effects on the occurrence and extent of atherosclerosis in rhesus monkeys. *J Clin Invest* 1980 Jan; 65:15-25
 Walker AM, Jick H, Hunter JR, et al: Vasectomy and non-fatal myocardial infarction. *Lancet* 1981 Jan 3; 1(8210):13-15

Computerized Tomography, Lymphangiography and Gleason Scores for Staging Prostatic Adenocarcinoma

IT HAS BEEN well documented that the prognosis and metastatic progression of disease in patients who have adenocarcinoma of the prostate is determined by the presence or absence of pelvic lymph node metastasis. There is no information that supports an increased survival with either pelvic lymph node dissection or pelvic lymph node irradiation. The accurate preopera-